Teagasc Dairy Beef Research Programme



Johnstown Castle, Co. Wexford

Update: July 2020

With the warm temperatures and rainfall, growth has remained high across the farm for the month of July, averaging 70 kg DM/ha. A number of paddocks, both trial and commercial, were cut as surplus bales on the 31^{st} July as they became too strong. The yearlings are performing well, gaining 0.90 kg/day, while the calves are gaining 0.77 kg/day. The remainder of the 2018 trial bullocks and 27 2018 spring born Friesian bullocks from the commercial herd were killed the end of July.

| | July 2020 |
|-----------------------|------------------|
| Total rainfall (ml) | 82.1 |
| Mean temperature (°C) | 14.6 |

Animal Performance:

28-month old steers

• The bullocks were killed on the 20th and 24th July. They averaged 670 kg live weight, and gained an average 1.05 kg/day during the third season at grass.

Table 1. Steer slaughter results, July 2020

| | Age (months) | LW (kg) | CW (kg) | KO% | Conf. | Conf. score | Fat. | Fat score |
|--------|-----------------|------------|---------|------|-------|----------------|------|--------------|
| High | 28.8 | 675 | 363.5 | 53.8 | 5.8 | O=/O+ | 9.6 | 3+/4- |
| Medium | 28.8 | 660 | 354.5 | 53.7 | 5.8 | O=/O+ | 9.5 | 3+/4- |
| Low | 28.6 | 680 | 362.5 | 53.3 | 5.5 | O=/O+ | 9.4 | 3+/4- |

Table 2. Total steer slaughter results March – July 2020

| | No. animals | Age (months) | LW (kg) | CW (kg) | KO% | Conf. | Conf. score | Fat. | Fat score |
|----|----------------|-----------------|------------|------------|------|-------|----------------|------|--------------|
| AA | 37 | 25.2 | 654 | 326 | 49.8 | 4.5 | O-/O= | 8.7 | 3=/3+ |
| HE | 36 | 25.5 | 636 | 317 | 49.8 | 4.6 | O-/O= | 9.6 | 3+/4- |
| LM | 27 | 27.8 | 667 | 352 | 52.7 | 6.2 | O+ | 9.2 | 3+ |

Yearlings (18-month-old)

- The high stocking rate yearlings' growth rate has picked up since the drought, now averaging 0.82 kg/day for the second season at grass.
- The yearlings were weighed on the 16th July 2020, and they received Spotinor pour-on for flies.
- The heifers and steers within the yearling groups have now been separated for grazing.
- The heifers will begin meal supplementation mid-August.

Table 3. Yearling weight and ADG

| | Live wei | ADG second season | | | |
|--------|----------|-------------------|----------|--|--|
| | Heifer | Steer | (kg/day) | | |
| High | 382 | 406 | 0.82 | | |
| Medium | 392 | 410 | 0.99 | | |
| Low | 398 | 409 | 0.88 | | |



Teagasc Dairy Beef Research Programme

Johnstown Castle, Co. Wexford

Spring 2020 calves

- The calves received pour-on (Spotinor) for flies on the 8th July.
- The calves were weighed and sorted into their treatment groups on the 2nd July. They averaged 113 kg and have been grazing their alloted treatments since.
- Following faecal samples, the calves were dosed for coccidiosis (Chanox) on the 22nd July. They also received their clostridia booster vaccination (Tribovax 10) and weighed 129 kg.
- The calves have been reduced to 0.42 kg concentrates/day.

Table 4. Calf live weight and ADG

| | LW 08/07/2020 | LW 22/07/2020 | ADG first season | ADG grazing trial plots |
|---------------------|------------------|------------------|---------------------|----------------------------|
| PRG | 116 | 128 | 0.75 | 0.78 |
| PRG + Clover | 118 | 133 | 0.84 | 0.88 |
| PRG + Clover + Herb | 111 | 128 | 0.72 | 0.85 |

Grassland Management

- Demand has reduced across all the stocking rate groups with the slaughter of the 28-month-old bullocks. Grass growth rate has averaged 70 kg DM/ha/day, which surpasses demand.
- The second cut silage was cut on the 10^{th} July and averaged 3715 kg DM/ha.
- Post second cut silage harvest, the silage ground was spread with 150 kg KAN and received 3000 gal/acre slurry before being closed on the 16th July 2020 with the aim to cut early September.
- On July 13th, a round of KAN was spread across the grazing paddocks at 65 kg/ha.
- Super P 16% was spread on the 20th July; rates of 45 to 125 kg/ha were applied dependent on the soil test results.
- Due to the reduction in demand and the increase in grass growth, a number of surplus paddocks, including some of the reseeded PRG and multispecies paddocks, were cut for surplus bales on the 31st July.
- The cattle have been grazing covers of 1950 kg DM/ha on average, grazing to heights of 4.39 cm.

| | Farm cover | Growth/ha | Demand /ha | Grazing area | Silage cut later | Total LU | LU/ha | Days ahead |
|------|---------------|-----------------|-----------------|-----------------|---------------------|-------------|-------|---------------|
| | Kg DM/ha | Kg DM/ha/day | Kg DM/ha/day | На | На | | | |
| High | 544 | 105 | 61 | 6.70 | 6.76 | 50.4 | 5.37 | 9 |
| Med | 620 | 93 | 58 | 8.53 | 7.67 | 50.4 | 5.04 | 11 |
| Low | 649 | 99 | 51 | 9.21 | 8.24 | 50.4 | 4.45 | 13 |

Table 5. Farm cover, 27th July 2020