

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Emerger (PCS No. 06190)

CROPS:

[Outdoor grown: parsnip, carrot, parsley root, onion, garlic, shallot, parsley & dill]

This approval provides for the use of the product **Emerger (PCS No. 06190)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 05.01.19. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Emerger
<i>Active Substance:</i>	Aclonifen
<i>Formulation:</i>	600g/l SC
<i>Authorisation Holder:</i>	Bayer CropScience Limited
<i>Marketing Company:</i>	Bayer CropScience Limited
<i>PCS No:</i>	06190
<i>User type/Function:</i>	Professional Use Only / Herbicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Parsnip	1.5 l/ha	2 per crop	2.5 l/ha	n/a	First application before germination, second at two true leaf stage and weeds in cotyledon stage	See Below*
Carrot, parsley root.	2.5 l/ha	2 per crop	2.5 l/ha	n/a	First application before germination, second at two true leaf stage and weeds in cotyledon stage	See Below*
Onion, garlic shallot	2.5 l/ha	2 per crop	2.5 l/ha	n/a	First application before germination, second at 1 to 2 true leaf stage and weeds in cotyledon stage	See Below*
Dill	2.5 to 3.0 l/ha	1 per crop	3.0 l/ha	n/a	After sowing and before germination	See below*

* Annual meadow grass, cleavers, common chickweed, fat hen, field pansy, henbit dead nettle, pale persicaria, red dead nettle, scented mayweed, shepherd's purse & volunteer oilseed rape.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 10m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Emerger as a horticultural herbicide for the control of: Annual meadow grass, cleavers, common chickweed, fat hen, field pansy, henbit dead nettle, pale persicaria, red dead nettle, scented mayweed, shepherd's purse & volunteer oilseed rape on the crops & situations listed above.
Contact your adviser for more info on water volumes etc.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, HRAC group] with satisfactory efficacy against the target.

Emerger contains Aclonifen

Aclonifen is classed HRAC group: F3 .

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.