

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Amistar (PCS No. 05072)

CROPS:

[Outdoor grown crops of: ornamentals and forest nurseries].

For this EAMU the following crop definitions apply:

Forest nurseries includes Christmas tree nurseries.

This approval provides for the use of the product **Amistar (PCS No. 05072)** in respect of, the crops and situations as listed above, which are additional to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 17.04.2018 [Includes latest changes. This version should be used until updated]*Product name:* Amistar

Active Substance: Azoxystrobin

Formulation: 250g/l SC

Authorisation Holder: Syngenta UK Ltd

Marketing Company: Syngenta Ireland Ltd

PCS No: 05072

User type/Function: Professional Use Only / Fungicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. single dose	Max. Number of applications	Max. Total dose	Minimum interval between applications	Latest time of application. PHI.	Target
[Outdoor grown crops of: Flowers, foliage, woody ornamentals, bulbs, ornamental seedlings] Forest nurseries.	0.7 l/ha	n/a	1.4 l/ha	7 days	-	leaf spots, powdery mildew & rusts.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed/air assisted or knapsack sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) The Pre harvest interval also applies to unprotected workers handling treated crops. That is for example: if the PHI is 2 days then unprotected workers may not handle treated material for 2 days post application.
- (6) Spraying may not occur at times when the public have any access to the vicinity of application.
- (7) The seed or plant material may not be used for human or animal feed.
- (8) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (9) This approval extends to professional use only.
- (10) For operator protection refer to the product label in conjunction with the specific restrictions above.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (11) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Amistar as a horticultural fungicide for the control of the listed: diseases on the crops and situations listed above.

Crop should not be under any stress at time of application. Do not use an adjuvant.

Do not apply at temperatures above 30C or below 10C. It is advisable to test a small area before use.

Amistar contains azoxystrobin, a broad spectrum fungicide from the strobilurin, QoI group. It has systemic, translaminar and protectant properties.

Azoxystrobin inhibits fungal respiration. It should always be used in mixture with fungicides with other modes of action. Amistar is best used as a protective treatment or during early stages of disease establishment.

Resistance Management:

This product should be applied in accordance with FRAC guidelines.

Disease control may be reduced if resistant strains of the pathogen less sensitive to azoxystrobin develop.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Amistar contains Azoxystrobin, which is classed FRAC Group 11, QoI fungicides.

Important please note:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.