

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Amistar (PCS No. 05072)

CROPS:

[Outdoor grown: Babyleaf crops, herbs, spinach, spinach beet, blackberry, raspberry, celery, cucumber, courgette, gherkin, melon, pumpkin, squash, marrow, Chinese cabbage, celeriac, swede, turnip, beetroot, horse radish, parsley root, parsnip, salsify & radish & salad onion]

[Indoor grown: Tomato, Aubergine, babyleaf crops, herbs, spinach, spinach beet, blackberry, raspberry, celery, cucumber, courgette, gherkin, melon, pumpkin, squash, marrow, pepper, chilli pepper,]

For this EAMU the following crop definitions apply:

Baby leaf crops include: baby leaf lettuce and baby leaf brassicas up to 8 true leaves, Butterhead, oakleaf, lollo rossa, lambs, iceberg, cos romaine, batavia, apollo, cress, land cress, rocket/wild rocket, red mustard, mizuna, pea shoots, salanova, red chard, tatsoi, bullsblood & multileaf types of the above list.

Perennial herbs: {Angelica, balm, bay, burnet (salad), catnip, chamomile, chives, clary sage, dragonhead, fennel, horehound, hyssop, land cress, lavandin, lavender, lemon verbena, lovage, marjoram, mint, nettle, oregano, plantain, rosemary, rue, sage, sorrel, spike lavender, St john's Wort, tarragon, thyme, woodruff, yarrow}

Annual herbs: {basil, Borage, caraway, chervil, coriander, curry plant, dill, fenugreek, feverfew, marigold, nasturtium, parsley, savory, violet}

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

This approval provides for the use of the product **Amistar (PCS No. 05072)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 17.04.18. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Amistar
<i>Active Substance:</i>	Azoxystrobin
<i>Formulation:</i>	250g/l SC
<i>Authorisation Holder:</i>	Syngenta UK Ltd
<i>Marketing Company:</i>	Syngenta Ireland Ltd
<i>PCS No:</i>	05072
<i>User type/Function:</i>	Professional Use Only / Fungicide

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
[Indoor tomato & aubergine]	1.0 l/ha	n/a	3.0 l/ha	3 days	3 days before harvest	Botrytis, powdery mildew & Didymella
[Indoor & outdoor; babyleaf crops, herbs, spinach, spinach beet]	1.0 l/ha	n/a	2.0 l/ha	14 days	14 days before harvest	Downy mildew, rhizoctonia
[Indoor & outdoor; blackberry & raspberry]	1.0 l/ha	n/a	2.0 l/ha	7 days	7 days before harvest	Powdery mildew
[Indoor & outdoor; celery]	1.0 l/ha	n/a	3.0 l/ha	14 days	14 days before harvest	Rhizoctonia, sclerotinia & botrytis
[Indoor & outdoor; Squash, cucumber, melon, courgette, gherkin, pumpkin, & marrow]	1.0 l/ha	n/a	3.0 l/ha	3 days	3 days before harvest	Powdery & downy mildew
[Indoor pepper & chilli pepper]	1.0 l/ha	n/a	2.0 l/ha	3 days	3 days before harvest	Powdery mildew
[Outdoor; Chinese cabbage]	1.0 l/ha	n/a	2.0 l/ha	14 days	14 days before harvest	White blister, ring spot & alternaria
[Outdoor; celeriac, swede, turnip, beetroot]	1.0 l/ha	n/a	2.0 l/ha	14 days	14 days	Alternaria, powdery mildew
[Outdoor; horse radish, parsley root, parsnip & salsify]	1.0 l/ha	n/a	3.0 l/ha	14 days	14 days before harvest	Alternaria, powdery mildew
[Outdoor; radish]	1.0 l/ha	n/a	2.0 l/ha	7 days	7 days before harvest	Alternaria & powdery mildew
[Outdoor; salad onion]	1.0 l/ha	n/a	3.0 l/ha	21 days	21 days before harvest	Downy mildew

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed/air assisted or knapsack sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label and specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies. Direct spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Amistar as a horticultural fungicide for use on the crops and situations listed above.

It is recommended to apply in 750 l/water/ha for blackberry and raspberry, 200 l/water/ha for celery and 300 l/water/ha for all the other crops.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in rotation & in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Amistar contains Azoxystrobin

Azoxystrobin is classed FRAC group: 11 QoI fungicides:

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

It is recommended that Amistar is used in an alternating programme where only one application in three is a QoI-based fungicide

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.