

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Biscaya (PCS No. 03602)

CROPS: Outdoor crops of: Beetroot, celeriac, horseradish, parsley root, salsify, swede, turnip, chinese cabbage, kale, collards, field beans.

This approval provides for the use of the product **Biscaya (PCS No. 03602)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 01.08.18. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Biscaya
<i>Active Substance:</i>	Thiacloprid
<i>Formulation:</i>	240 g/l Oil dispersion formulation
<i>Authorisation Holder:</i>	Bayer CropScience Ltd.
<i>Marketing Company:</i>	Bayer CropScience Ltd.
<i>PCS No:</i>	03602
<i>User type/Function:</i>	Professional Use Only / Insecticide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Beetroot, horseradish, parsley root, salsify,	0.4 l/ha	2 per crop	0.8 l/ha		14 days before harvest	Aphids, carrot willow aphid
Celeriac	0.4 l/ha	2 per crop	0.8 l/ha		7 days before harvest	Aphids, carrot willow aphid, peach potato aphid, black bean aphid
Swede & turnip	0.4 l/ha	2 per crop	0.8 l/ha		35 days before harvest	Aphids, mealy cabbage aphid
Kale, collards	0.4 l/ha	2 per crop	0.8 l/ha		21 days before harvest	Mealy cabbage aphid
Chinese cabbage including choi sum, Pak Choi, tat soi	0.4 l/ha	2 per crop	0.8 l/ha		7 days before harvest	Mealy cabbage aphid
Field bean *	0.4 l/ha	2 per crop	0.8 l/ha	10 days	7 days before harvest	Pea aphid, bruchid beetle

* Applications to field bean must be made between 1st May and 31st July only.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a tractor mounted/trailed hydraulic sprayer only.
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (5) This approval extends to professional use only.
- (6) For operator protection refer to the product label.
- (7) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Biscaya as an agricultural insecticide for the crops & situations listed above.

Resistance Management:

Use Biscaya as part of an Integrated Crop Management (ICM) strategy.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other insecticides from different cross-resistant groups [different mode of action, IRAC group] with satisfactory efficacy against the target.

Biscaya contains Thiacloprid IRAC class: 4A.

Kale, collards & Chinese cabbage: Use no more than two foliar applications of any neonicotinoid insecticide (eg. clothianidin, imidacloprid, thiacloprid, acetamiprid) on any crop. Two foliar applications are permitted regardless of whether a neonicotinoid insecticide has previously been used on the crop as a seed or module treatment.

All other crops: Do not use more than two applications of any neonicotinoid insecticide (e.g. acetamiprid, clothianidin, imidacloprid or thiacloprid) either foliar or seed treatment on any crop per year. Alternate with products with different modes of action. Where a neonicotinoid insecticide has previously been used as a soil treatment, use no more than a single foliar application of any neonicotinoid.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.