

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Movento (PCS No. 05349)

CROPS: Parsnip (outdoor), swede (outdoor), turnip (outdoor), shallot (outdoor), chinese cabbage (outdoor), kohlrabi (outdoor), baby leaf production (outdoor & protected), Herbs_(outdoor & protected), spinach beet (outdoor & protected), endives (outdoor & protected), spinach (outdoor), chicory (outdoor). Protected crops of tomato, aubergine, cucumber, pepper, melon, and courgette.

This approval provides for the use of the product **Movento (PCS No. 05349)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 05.11.18. [Includes latest changes. This version should be used until updated]

Product name: Movento

Active Substance: Spirotetramat

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Formulation: 150g/l Oil dispersion formulation

Authorisation Holder: Bayer CropScience Ltd.

Marketing Company: Bayer CropScience Ltd.

PCS No: 05349

User type/Function: Professional Use Only / Insecticide.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Outdoor crops of chinese cabbage, kohlrabi	0.5 L/ha	2 per crop	1.0 L/ha	14 days	3 days before harvest	Aphids, whitefly, mealy cabbage aphid (<i>Brevicoryne brassicae</i>), peach potato aphid (<i>Myzus persicae</i>), Lettuce root aphid (<i>Pemphigus bursarius</i>) and Brassica whitefly (<i>Aleyrodes proletella</i>)

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Protected and outdoor crops of baby leaf production, herbs, spinach beet, endive	0.5 L/ha	2 per crop	1.0 L/ha	14 days	7 days before harvest	Aphids, whitefly, mealy cabbage aphid (<i>Brevicoryne brassicae</i>), peach potato aphid (<i>Myzus persicae</i>), Lettuce root aphid (<i>Pemphigus bursarius</i>) and Brassica whitefly (<i>Aleyrodes proletella</i>)
Outdoor crops of spinach, chicory	0.5 L/ha	2 per crop	1.0 L/ha	14 days	21 days before harvest	Aphids, lettuce root aphid
Parsnip	0.5 L/ha	2 per crop	1.0 L/ha	14 days	21 days before harvest	Carrot psyllid
Swede & Turnip	0.5 l/ha	2 per crop	1.0 l/ha	14 days	21 days before harvest	Mealy cabbage aphid, peach potato aphid
Outdoor shallot	0.5 l/ha	4 per crop	2.0 l/ha	14 days	7 days before harvest	Onion thrip
Protected crops of tomato, aubergine, cucumber, melon, pepper, courgette	0.5 l/ha	4 per crop	2.0 l/ha	14 days	14 days before harvest	Aphids, greenhouse whitefly, silverleaf whitefly (<i>Trialeurodes vaporariorum</i> , <i>Bemisia tabaci</i>), Western flower thrips

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Dangerous to bees. To protect bees and other pollinating insects do not apply to crop plants when in flower or when flowering weeds are present. Do not apply when bees are actively foraging.
- (4) Application may only be made using a tractor mounted/trailed hydraulic sprayer.
- (5) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (6) Apply in a water volume of 200-600 l/ha. For protected tomato, cucumber, aubergine, melon, pepper and courgette apply in a minimum of 1500 l/ha
- (7) This approval extends to professional use only.
- (8) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water.
- (9) For operator protection refer to the product label.
- (10) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Movento as a horticultural insecticide for the control of a range of pests as prescribed on the crops & situations listed above.

Resistance Management:

Use Movento as part of an Integrated Crop Management (ICM) strategy. In a spray programme Movento should be used with other insecticides of a different mode of action, either in alternation or as a 2-spray block within the programme

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy.

Movento contains Spirotetramat IRAC class: 23 which also includes spirodiclofen and spiromesifen.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.